

CHERRY

See also the general sheet **FRUIT GROWING**

PERIOD	PRODUCT	DOSE	Application	NOTES
Pre-flowering	Alga Special + Boron 21 PG	1.5 l/ha 1 kg/ha	Foliar application	Improves setting
At beginning of flowering	Alga Special + Boron 21 PG	1.5 l/ha 1 kg/ha	Foliar application	Improves size
After 4 days	Alga Special	3 l/ha	Foliar application	Improves size and prevents May drop.
Post-flowering	Alga Special +Calcium 44 LG + Sprühdünger type 5	3.5 l/ha 750 ml/ha 750 g/ha	Foliar application	Prevents fruit splitting
After setting and up until veraison	Alga Special + Sprühdünger type 5	3.5 l/ha 750 g/ha	Foliar application	Deliver at least three treatments at 7-8 day intervals.
Upon veraison	Sprühdünger type 30 +Alga Special +Calcium 44 LG	1 kg/ha 2 l/ha 1 l/ha	Foliar application	Prevents fruit splitting

Alternative to blend

Alga Special + Boron 21 PG	1.5 l/ha 1 kg/ha
---	-----------------------------------

Can be used

Alga Special + Boro LG 25	1.5 l/ha 2.5 l/ha
--	------------------------------------

Alga Special + Boro 10 LG	1.5 l/ha 1 l/ha
--	----------------------------------

or

Alga Mix BMg	3 l/ha
---------------------	---------------

The new product **SEAWEED MIX** can be used in any application for which the use of **ALGA SPECIAL** is recommended: it boasts a very high concentration obtained from ASCOPHYLLUM NODOSUM extracts and from LAMINARIA DIGITATA concentrate and cream.

in case of need, as an alternative to **SPRÜHDÜNGER type 5** one can use **SPRÜHDÜNGER type 30** which includes a higher potassium contents

As an alternative to **ALGA SPECIAL + CALCIUM 44 LG**, **ALGA CA** dosed at 3 l/ha



I- 16013 Campo Ligure (GE) Via Vallecaldà, 33
Tel.: +39 010 920 395 Fax : +39 010 921 400
e-mail: info@lgobbi.it http://www.lgobbi.it

SPECIALITÀ PER L'AGRICOLTURA

l. gobbi s.r.l.

Conc. E. Gerlach GmbH - Germania

