

# COURGETTE

SEE THE GENERAL SHEET " HORTICULTURE "

PERIOD	PRODUCT	DOSE	Application	NOTA
TRANSPLANT TO BED	<b>Alga Special</b>	<b>2 l/1000 m<sup>2</sup></b>	Fertirrigation	Promotes a rapid recovery and helps seedlings live through transplant shock.
	<b>Stimolante 66f + Gerfos-K</b>	<b>20 ml/hl 300 ml/hl</b>	Foliar application	Repeat every 15-20 days
IN THE PERIOD THAT FOLLOWS	<b>Stimolante 66f + Sprühdünger type 27 + Gerhumin</b>	<b>20 ml/hl 50 g/hl 50 ml/hl</b>	Foliar application	Promotes quick and balanced growth of plants. Apply weekly.
FLOWERING	<b>Adrop + Alga Special</b>	<b>50 – 100 g/hl 100 ml/hl</b>	Foliar application	Promotes setting. Apply during flowering, every 8-10 days, progressively increasing ADROP dose (do not exceed 100 g/hl).
AFTER 4-5 DAYS	<b>Stimolante 66f + Alga Special + Sprühdünger type 26 + Boro LG 25</b>	<b>20 ml/hl 100 ml/ hl 50 g/hl 400 ml/hl</b>	Foliar application	
	<b>Sprühdünger type 26 + Boro LG 25 + Gerhumin</b>	<b>2 kg/1000 m<sup>2</sup> 4 l/1000 m<sup>2</sup> 1 l/1000 m<sup>2</sup></b>	Radical application	
SETTING / BEGINNING OF RIPENING	<b>36C Type A (or Adrop or Gerauxin) + Seaweed Mix + Sprühdünger type 5</b>	<b>100 ml/hl 50 g/hl 30-50 ml/hl 200 ml/hl 50 g/hl</b>	Foliar application	Promotes fruit development and a smoother and more balanced ripening.
	<b>Fluidünger Più + Gerhumin</b>	<b>10 l/1000 m<sup>2</sup> 1 l/1000 m<sup>2</sup></b>	Radical application	
BEGINNING OF HARVESTING	<b>Sprühdünger type 5 + Gerhumin</b>	<b>100 g/hl 50 ml/hl</b>	Foliar application	Covers potassium requirements of courgette at harvesting stage. A deficiency would seriously affect yield, in terms of quality and quantity.
	<b>Fluidünger Più + Gerhumin</b>	<b>10 l/1000 m<sup>2</sup> 1 l/1000 m<sup>2</sup></b>	Radical application	



I- 16013 Campo Ligure (GE) Via Vallecaldà, 33  
Tel.: +39 010 920 395 Fax : +39 010 921 400  
e-mail: info@lgobbi.it <http://www.lgobbi.it>

SPECIALITÀ PER L'AGRICOLTURA

**l. gobbi** s.r.l.

Conc. E. Gerlach GmbH - Germania

