

# PEAR

See also the general sheet **FRUIT GROWING**

PERIOD	PRODUCT	DOSE	Application	NOTES
Early Spring	<b>LG 81</b> <b>+ Anticlor Fe WP6G</b>	<b>6 l/ha</b> <b>30-50 g/tree</b>	To the soil	It improves the soil and of the roots biological activity increasing the calcium uptake.
Tight clusters	<b>Stimolante 66f</b> <b>+ Alga CMZ</b> <b>+ Boron 21 PG</b>	<b>300 ml/ha</b> <b>3 l/ha</b> <b>1 kg/ha</b>	Foliar application	Protection of flowering against adverse weather conditions
Loose clusters	<b>Stimolante 66f</b> <b>+ Alga CMZ</b> <b>+ Boron 21 PG</b> <b>+ Sprühdünger 26</b>	<b>300 ml/ha</b> <b>3 l/ha</b> <b>1 kg/ha</b> <b>750 g/ha</b>	Foliar application	Preparation for an optimized setting
10-20% of open flowers	<b>AF 96</b> <b>+ Alga Mix BMg</b> <b>+ Sprühdünger 26</b>	<b>800-1200 ml/ha</b> <b>3 l/ha</b> <b>750 g/ha</b>	Foliar application	Increase in setting and fruit formation, according to variety.
<b>Abate Fétèl</b>	Treat with <b>AF 96</b> when approx. 10-20% of flowers are open, then repeat when 50% are open. Alternative treatment: apply <b>AF 96</b> (1 l/ha) on appearance of white buds, then treat with <b>Progerbalin</b> (200-300 ml/ha) when 10-20% of flowers are open, then repeat using the same dose after 2-3 days.			
or: 25% of open flowers	<b>Progerbalin</b>	<b>60-80 ml/hl</b> (or 30-40 ml/hl in case of two treatments)	Foliar application	Increase in setting. Apply a second time 2-3 days after the first.
or: 30-40% of open flowers	<b>Gerlagib LG</b>	<b>80-100 ml/hl</b> (or 30-50 ml/hl in case of two treatments)	Foliar application	
On petal fall	<b>Stimolante 66f</b> <b>+ Sprühdünger 26</b> <b>+ Alga Mix BMg</b>	<b>300 ml/ha</b> <b>750 g/ha</b> <b>3 l/ha</b>	Foliar application	Fruit formation
2-3 weeks after full bloom	<b>Platina</b>	<b>1.5 l/ha</b>	Foliar application	Improved growth rate and fruit development
Mid-May	<b>Alga Special</b> <b>+ Sprühdünger 26</b>	<b>2 l/ha</b> <b>750 g/ha</b>	Foliar application	
End of May	<b>Sprühdünger 26</b>	<b>750 g/ha</b>	Foliar application	
Beginning of June	<b>Alga Special</b> <b>+ Sprühdünger 26</b>	<b>2 l/ha</b> <b>750 g/ha</b>	Foliar application	
Mid-June End of June Beginning of July	<b>Alga Special</b> <b>+ Sprühdünger 5</b> <b>+ Zinc 10 LG</b>	<b>3 l/ha</b> <b>750 g/ha</b> <b>1.5 kg/ha</b>	Foliar application	Better colouring and increase in pulp consistency, improved shelf live and augmented differentiation of flower buds.
End of July	<b>Alga Special</b> <b>+ Sprühdünger 5</b>	<b>3 l/ha</b> <b>750 g/ha</b>	Foliar application	Every fifteen days up until a few days before harvesting, so as to improve fruit storability.
7-20 days prior to harvest	<b>Obsthormon 24a</b>	<b>250-500 ml/ha</b> according to variety	Foliar application	Prevention of pre-harvesting drop.
Soon after harvesting	<b>Alga Special</b> <b>+ Boron 21 PG</b> <b>+ Urea</b>	<b>(3 l/ha)</b> <b>(1 kg/ha)</b>	Foliar application	It stimulates accumulation to ensure maximum availability during the early stages of the following season.
Autumn	<b>LG 81</b>	<b>6 l/ha</b>	To the soil	It improves the soil and of the roots biological activity increasing nutrient uptake.

Notice to the listed products:

**Boron 21 PG** (1 kg/ha) can be replaced by **Boro LG 25** (2,5 l/ha) or by **Boro 10 LG** (1 l/ha).

The blend **Alga Special** (2-3 l/ha) + **Boron 21 PG** (1 kg/ha) can be replaced by **Alga Mix BMg** (3 l/ha).

**Alga Special** can be replaced by **Seaweed Mix** (same doses).



I- 16013 Campo Ligure (GE) Via Vallecaldà, 33  
Tel.: +39 010 920 395 Fax : +39 010 921 400  
e-mail: info@lgobbi.it http://www.lgobbi.it

**SPECIALITÀ PER L'AGRICOLTURA**

**l. gobbi** s.r.l.

Conc. E. Gerlach GmbH - Germania

