

Open field STRAWBERRY

SEE THE GENERAL SHEET " HORTICULTURE "

PERIOD	PRODUCT	DOSE	Application	NOTES
PRIOR TO TRANSPLANT	Stimolante 66f	20 ml/hl	Root dipping	Prevents transplant shock. Dip seedlings for approx. 12 hours. Also mixed with treatments for the prevention of root rotting. For cold stored seedlings, it is recommended to take them out of the store room at least 1-2 days before treatment.
AFTER TRANSPLANTING	Stimolante 66f + Gerfos-K	20 ml/hl 50 ml/hl	Foliar application	Repeat every 15-20 days.
	Alga Special	150 ml/hl	Radical application	Promotes immediate and uniform plant growth and creates the ideal conditions for a profuse flowering.
ON VEGETATIVE RECOVERY	Anticlor Fe WP6 G + Alga Special	100 g/hl 150 ml/hl	Radical application	For a couple of times
AFTER SOME DAYS	Alga Special + Sprühdünger type 26	150 ml/hl 50 g/hl	Radical application	For a couple of times
POST-FLOWERING (Beginning of petal fall)	36C Type A (or Adrop Polvere or Gerauxin) +Alga Special	100 ml/hl 50 g/hl (30-50 ml/hl) 150 ml/hl	Foliar application	Promotes setting.
AFTER SETTING	Stimolante 66f + Alga Special + Sprühdünger type 5	20 ml/hl 150 ml/hl 50 g/hl	Foliar application	Promotes a quick and uniform growth rate of the small fruit, positively influencing its organoleptic characteristics.
	Alga Special +Sprühdünger type 5	150 ml/hl 50 g/hl	Radical application	

Deficiencies: Calcium

PERIOD	Alternative 1		Alternative 2		Alternative 3		
During fruit growth. (Early growth stages)	Alga Special	500 ml/hl	Alga Special	500 ml/hl	Alga Ca	2 l/hl	Radical application
	TRB 124	1.5 kg/hl or	+ Calcium 44 LG	300-400 g/hl or			
During fruit growth. (Close to ripening)	Alga Special	150 ml/hl	Alga Special	150 ml/hl	Alga Ca	500 ml/hl	Foliar application
	TRB 124	400 g/hl or	+ Calcium 44 LG	80-100 g/hl or			



I- 16013 Campo Ligure (GE) Via Vallecaldà, 33
Tel.: +39 010 920 395 Fax : +39 010 921 400
e-mail: info@lgobbi.it http://www.lgobbi.it

SPECIALITÀ PER L'AGRICOLTURA

l. gobbi s.r.l.

Conc. E. Gerlach GmbH - Germania

