



# SUGAR BEET

## - RECOMMENDED SCHEDULE FOR SUGAR CONTENT INCREASE AND OPTIMAL ROOTING PERFORMANCE

VEGETATIVE STAGE	PRODUCT	DOSES	INDICATIONS
From 4 leaves → Rosette 	Stimolante 66f	0.3 l/ha	Apply at least twice, mixed with post-emergency herbicide treatments
taproot development 	Stimolante 66f + Sprühdünger type 5	0.3 l/ha + 0.75 kg/ha	Apply once or twice after 10-15 days, at the same time as any cercospora fungicide treatment

## - CHECK OF MAIN DEFICIENCY AND DISEASE -

Apply upon appearance of the first symptoms with at least 2 treatments at 10 – 15 days' interval

DEFICIENCY - DISEASE	PRODUCT	DOSES	INDICATIONS
<b>BORON</b>	Boro 10 LG	1.5 - 2 kg/ha	As this deficiency is quite common, we recommend that you carry out systemic applications, associated to the previous Program
<i>Manganese</i>	Mangan 32 PG	1 kg/ha	
<i>Magnesium</i>	Magnesium 16 PG	5 - 8 Kg /ha	
<i>Multiple deficiencies</i>	EG 118	3 – 4 kg/ha	Use this formula if more than one nutrient is deficient and/or if deficiency cannot be related to a single element.
<i>Nitrogen</i>	Sprühdünger type 27 (*)	750 g/ha	(*) After 6-8 leaves, Sprühdünger type 5 is to be preferred
<i>Poorly structured soil</i> (timely tillage, stagnant water, etc.)	Stimolante 66f + Sprühdünger type 27 (*)	300 ml/ha + 750 g/ha	Leaf yellowing similar to nitrogen deficiency. (*) After 6-8 leaves, Sprühdünger type 5 is to be preferred

In deficiency treatment, a mixture with ALGA SPECIAL is recommended (2-3 l/ha), if you are not already using Stimolante 66f.



I- 16013 Campo Ligure (GE) Via Vallecaldà, 33  
 Tel.: +39 010 920 395 Fax : +39 010 921 400  
 e-mail: info@lgobbi.it http://www.lgobbi.it

SPECIALITÀ PER L'AGRICOLTURA

**l. gobbi** s.r.l.

Conc. E. Gerlach GmbH - Germania



## SEED CROPS

<i>VEGETATIVE STAGE</i>	<i>INSTRUCTIONS FOR USE</i>	<i>PRODUCT</i>	<i>DOSES</i>	<i>PURPOSE</i>
<i>Pre-transplanting</i>	Dip the root system into the solution or apply to the whole plant using a sprayer	<b>Stimolante 66f</b>	<b>20 ml/hl</b>	Improves post-transplanting performance
<i>Durante the whole growth stage</i>	Apply 3 - 4 times ever 15 – 20 days	<b>Stimolante 66f + Dünger 20</b>	<b>20 ml/hl + 150 g/hl</b>	Promote a vigorous vegetative development and a quick transition to the reproductive phase