

COURGETTE

SEE THE GENERAL SHEET "HORTICULTURE"

PERIOD	PRODUCT	DOSE	Application	NOTA
TRANSPLANT TO BED	Alga Special	2 l/1000 m²	Fertigation	Promotes a rapid recovery and helps seedlings live through transplant shock.
	Stimolante 66f + Gerfos-K	20 ml/hl 300 ml/hl	Foliar application	Repeat every 15-20 days
IN THE PERIOD THAT FOLLOWS	Stimolante 66f + Sprühdünger type 27 + Gerhumin	20 ml/hl 50 g/hl 50 ml/hl	Foliar application	Promotes quick and balanced growth of plants. Apply weekly.
FLOWERING	Auxiger + Alga Special	50 ml/hl 100 ml/hl	Foliar application	Promotes setting. Apply during flowering, every 8-10 days.
AFTER 4-5 DAYS	Alga Special + Sprühdünger type 26 + Boro LG 25	100 ml/ hl 50 g/hl 400 ml/hl	Foliar application	
	Sprühdünger type 26 + Boro LG 25 + Gerhumin	2 kg/1000 m² 4 l/1000 m² 1 l/1000 m²	Radical application	
SETTING / BEGINNING OF RIPENING	Auxiger + Seaweed Mix + Sprühdünger type 5	50 ml/hl 200 ml/hl 50 g/hl	Foliar application	Promotes fruit development and a smoother and more balanced ripening.
BEGINNING OF HARVESTING	Sprühdünger type 5 + Gerhumin	100 g/hl 50 ml/hl	Foliar application	Covers potassium requirements of courgette at harvesting stage. A deficiency would seriously affect yield, in terms of quality and quantity.

Use plant protection products with caution. Before use always read the label and the product information.
Pay attention to the hazard sentences and symbols shown on the label.



l.gobbi srl
unipersonale

Via Vallecaldà, 33 - 16013 Campo Ligure (GE) - ITALY
Tel.: +39 010 920 395 - Fax: +39 010 921 400
www.lgobbi.it
Conc. E. Gerlach GmbH - Germany