


# Nutritional advises

The best L. Gobbi s.r.l.  
products advises

# Fruit growing



## Resolution of specific problems

### 1. Setting

In general, the following treatments are recommended in order to prepare the plant for an optimal setting:

TIMING	PRODUCT	DOSE	VIA	NOTES
Beginning-Full bloom (0-10% and 20-50%)	<b>Auxiger Alga Mix BMg</b>	<b>500-700 ml/ha 3 L/ha</b>	Foliar	2 treatments

### 2. Fruit growth

For the purpose of promoting fruit growth rate, the following treatments are generally recommended:

TIMING	PRODUCT	DOSE	VIA	NOTES
From setting to ripening	<b>Alga Special (or Seaweed Mix)</b>	<b>2 L/ha (2 L/ha)</b>	Foliar application	Fortnightly

Alternatively, in the crops where it is allowed:

TIMING	PRODUCT	DOSE	VIA	NOTES
From the end of petal fall and all over the period during cells division, fortnightly	<b>Auxiger + Alga Special (or Seaweed Mix)</b>	<b>500-700 ml/ha 2 L/ha (2 L/ha)</b>	Foliar application	To increase fruits size.

### 3. Frost

To reduce damage caused by a sudden frost, the following treatments are recommended:

TIMING	PRODUCT	DOSE	VIA	NOTES
Some days before the predicted event.	<b>CropAid AntiFrost</b>	<b>500 ml/hl</b>	Foliar application	Ideal should be to start the treatments some week before but the efficacy can be showed even 6-12 hours before the frost peak. Repeat further twice.
When frost is to be expected (Preventive treatment)	<b>Auxiger + Alga Mix BMg</b>	<b>500-700 ml/ha 3 L/ha</b>	Foliar application	When the frost period go on for a long time, the treatment should be repeated to ensure a longer protection.
In case of late or unexpected frost (treatment after the frost)	<b>Auxiger + Alga Mix BMg</b>	<b>500-700 ml/ha 3 L/ha</b>	Foliar application	Apply immediately within the next 24 hours

### 4. Suckers and side shoots

In general, in order to eliminate sucker and side shoots and reduce their emergence, the following treatments are recommended:

TIMING	PRODUCT	DOSE	VIA	NOTES
When the suckers are 10-15 cm in height.	<b>Spollonante G</b>	<b>1 l/10l</b> (the quantity per hectare varies on the basis of the number and development of suckers and on hectare culture intensity)	Foliar application	Mainly spray the sucker tip

Use plant protection products with caution. Before use always read the label and the product information.  
Pay attention to the hazard sentences and the symbols shown on the label.



**l.gobbi srl**  
unipersonale

Via Vallecaldà, 33 - 16013 Campo Ligure (GE) - ITALY  
Tel.: +39 010 920 395 - Fax: +39 010 921 400  
www.lgobbi.it  
Conc. E. Gerlach GmbH - Germany