



The L. Gobbi s.r.l products best advises

TIMING	PRODUCT	DOSE	VIA	NOTES
Early springtime	LG 81	4-6 L/ha	 Fertigation	It improves biological activity of soil and roots. Increases calcium absorption and uptake.
Every month (or every 15 days at half dosage)	LG 81	1 L/ha	 In fertigation along the raw	
Or at Midsummer		4-6 L/ha	 All cultivated soil surface	
Budding (Shoot 10 cm long)	Anticlor Fe WP6 G + Gerhumin	40-60 g/plant 30 ml/hl	 Fertigation	To avoid the iron deficiency and to enable plant to the flowering.
	Alga Special + Sprühdünger 27	2 l/ha 750 g/ha	 Foliar	
Pre/Beginning of flowering	Alga BFeMn	2 L/ha	 Foliar	To improve fruit setting. Repeat each treatment at least twice.
Beginning of flowering /Petals fall	Alga Special + Sprühdünger 26	2 L/ha 750 g/ha	 Foliar	
10 days after full bloom	Alga Ca	3 L/ha	 Foliar	To increase the fruit organoleptic features, increasing sugar content, fruit size and shelf life.
25 days after full bloom	Alga Ca	3 L/ha	 Foliar	
Beginning of fruit growing	Alga Special + Sprühdünger 26	2 L/ha 750 g/ha	 Foliar	Repeat the treatment at least twice. Keep 10 days one to the other.
40 days after full bloom	Alga Ca	3 L/ha	 Foliar	To increase the fruit organoleptic features, increasing sugar content, fruit size and shelf life.
Fruit set occurred	Alga Special + Sprühdünger 5 (o Sprühdünger 30)	2 L/ha 750 g/ha (750 g/ha)	 Foliar	Fortnightly
After the harvest	Alga Special + Energemma	3 L/ha 2 kg/ha	 Foliar	Si favorisce l'accumulo per una pronta disponibilità nelle primissime fasi della stagione successiva
20-30 days before the expected date of the leaves falling.	Blattab	8 kg/ha	 Foliar	To induce a regular leaves falling, sugar and nutrients transferring in the wood. It anticipates and increases the productivity in the following season! The treatment induces a better resistance to the frost. Apply 10 hl/ha of water.
10-14 days after.		8 kg/ha	 Foliar	
Autumn, after the leaves falling.	LG 81	4-6 L/ha	 Fertigation	It improves the soil and roots biological activity. It even improves the nutrients uptake and metabolism.
Control of the excessive vegetative emissions.	Brafix	q.b.	 Brushed	Apply with a spatula or a brush on large cuts and every time it was necessary to regulate the excessive vegetative development.

Use plant protection products safely. Before use always read the label and product information. Pay attention to the phrases and hazard symbols on the label.

Treatment with **Blattab** promotes leaf fall, wood maturation and increases plant resistance to frosts. Consequently, when the first frosts will occur, the plants, now completely at rest, will suffer less injuries, reducing the possibility of **PSA** infections, also due to the copper content of the product.



l.gobbi srl

Via Vallecaldà, 33 - 16013 Campo Ligure (GE) - ITALY

Tel.: +39 010 920 395 - Fax: +39 010 921 400

E-mail: info@lgobbi.it - <http://www.lgobbi.it>

Conc. E. Gerlach GmbH - Germany